

Cumulative Frequency Graphs


(a)

The heights of 40 oak trees are recorded in a frequency table.

(a) Use the information to calculate cumulative frequencies and plot a cumulative frequency graph.

(b) Use your cumulative frequency graph to find the median and interquartile range of the heights of the oak trees.

Height (m)	Frequency
$10 < h \leq 15$	5
$15 < h \leq 20$	9
$20 < h \leq 25$	13
$25 < h \leq 30$	7
$30 < h \leq 35$	4
$35 < h \leq 40$	2



(b)

The time taken by 100 students to complete a jigsaw is recorded in a grouped frequency table.

(a) Use the information to calculate cumulative frequencies and plot a cumulative frequency graph.

(c) Use your cumulative frequency graph to find the median and interquartile range of the times taken.

Time (minutes)	Frequency
$30 < t \leq 40$	4
$40 < t \leq 50$	11
$50 < t \leq 60$	20
$60 < t \leq 70$	28
$70 < t \leq 80$	19
$80 < t \leq 90$	13
$90 < t \leq 100$	5

(c)

The History test score of 60 students is recorded in a grouped frequency table.

(a) Plot a cumulative frequency graph.

(b) Find the median score in the History test.

(c) Use your graph to estimate the number of students who scored less than 36 marks.

(d) Use your graph to find the minimum score required to come in the top 10% of students in the test.


Test Score	Frequency
$0 < s \leq 10$	1
$10 < s \leq 20$	4
$20 < s \leq 30$	6
$30 < s \leq 40$	12
$40 < s \leq 50$	17
$50 < s \leq 60$	13
$60 < s \leq 70$	5
$70 < s \leq 80$	2

(b) Time Cumulative
Freq.

$30 < t \leq 40$	4
$40 < t \leq 50$	15
$50 < t \leq 60$	35
$60 < t \leq 70$	63
$70 < t \leq 80$	82
$80 < t \leq 90$	95
$90 < t \leq 100$	100

Median = 66 mins

IQR = 76 - 55
= 21 mins



(c) Test Score Cumulative Freq.

$0 < S \leq 10$	1
$10 < S \leq 20$	5
$20 < S \leq 30$	11
$30 < S \leq 40$	23
$40 < S \leq 50$	40
$50 < S \leq 60$	53
$60 < S \leq 70$	58
$70 < S \leq 80$	60

Median = 44 marks

< 36 marks

= 17 students

Top 10% score

= 62 marks.

